

is the 1,3,5-tribenzo derivative. Its formation requires extensive rearrangement, but it can be envisioned as arising from **7** by a rearrangement involving bond a, followed by a further bond rearrangement. Cleavage of bond b, however, produces **1**. Since the formation of **7** itself is unlikely, **1** being able to be formed instead without disruption of an aromatic system, the absence of a COT derivative is readily rationalized.

Thomas D. Walsh

Department of Chemistry, University of South Dakota
Vermillion, South Dakota 57069

Received October 21, 1968


Photochemistry of Triptycene¹

Sir:

Interest in an extension of previous work on the photochemistry² and the intriguing possibilities of excitation migration³ of benzobarrelenes prompted us to report our investigation of the photochemistry of some triptycenes.

Irradiation⁴ of triptycene (**1**) results in the smooth formation ($\Phi = 0.3$) of one product, **3**: mp 135.5–136.0°; nmr (CS₂, TMS internal reference) δ 1.25 (1 H, m), 2.40 (1 H, m), 4.46 (1 H, m), 6.10 (3 H, m), and 7.2–7.9 (8 H, m). Upon heating **3** is converted to 2,3-benzfluoanthene (**5**).⁵ Ozonolysis of **3** yields the keto aldehyde **4**: nmr (CS₂, TMS) δ 2.08 (1 H, d of d, $J = 6, 4$ cps), 3.28 (1 H, d of d, $J = 4, 1.5$ cps), 4.25 (1 H, d, $J = 1.5$ cps), 7.1–7.9 (8 H, m), 9.84 (1 H, d, $J = 6$ cps); mass spectra, m/e 260 (m^+ , 35%); ir max (CCl₄) 1709 and 1716 cm⁻¹. Compound **4** was converted to a diol by lithium aluminum hydride. Nmr spectra of this diol analysis demonstrated the presence of a tertiary benzylic (δ 5.6) and primary (δ 4.0) alcohol, thereby corroborating the structural assignment of **4**.

These results contrast with previous results on the photochemistry of barrelene and its benzo derivatives which yield a mixture of semibullvalenes and cyclo-octatetraenes when irradiated.² Formation of **3** can be interpreted in terms of formation of the semibullvalene **2** which then undergoes a sigmatropic rear-



angement⁶ to **3**. Models indicate that the rearrangement of **1** to **2** can plausibly lead only to the stereoisomer shown which likewise by sigmatropic rearrangement is most likely to yield stereoisomer **3**. The structural assignment is consistent with the low coupling constants⁷ between the cyclopropyl hydrogens of **3** ($J \sim 3$ cps) and **4** ($J = 4$ cps). The formation of **3** cannot be quenched by 1,3-pentadiene, but conversion of **1** to **3** also occurs smoothly in photosensitizing solvents such as benzene and acetone. Thus, the conversion of **1** to **3** may be a very fast triplet rearrangement, in analogy to those found for other barrelene derivatives. However, since formation of **3** is also observed in photolyses at 77°K (under conditions such that the phosphorescence ($\tau_p = 3$ sec) of **1** is observable), unless the rate of this rearrangement possesses a remarkably large temperature coefficient, it appears unlikely that the reaction occurs from T₁.

Compound **3** represents one of the few examples of a (nonbenzo) norcaradiene which is not the minor component in equilibrium with corresponding tropyliene⁸ at room temperature.

(1) (a) Molecular Photochemistry. XII. The authors at Columbia gratefully acknowledge the generous support of this work by the Air Force Office of Scientific Research (Grant AFOSR-68-1381). The authors at Case Western Reserve acknowledge the support from the National Cancer Institute (Public Health Service Research Grant No. CA-07909-03). (b) Presented in part at 156th National Meeting of the American Chemical Society, Atlantic City, N. J., Sept 1968, Abstracts No. PHYS-167.

(2) (a) P. W. Rabideau, J. B. Hamilton, and L. Friedman, *J. Am. Chem. Soc.*, **90**, 4465 (1968); (b) H. E. Zimmerman, R. S. Givens, and R. M. Pagni, *ibid.*, **90**, 4191 (1968); (c) H. E. Zimmerman, R. W. Binkley, R. S. Givens, and M. A. Sherwin, *ibid.*, **89**, 3932 (1967); (d) J. P. N. Brewer and H. Heaney, *Chem. Commun.*, 811 (1967); (e) R. S. H. Liu, *J. Am. Chem. Soc.*, **90**, 215 (1968); (f) E. Ciganek, *ibid.*, **88**, 2882 (1966).

(3) M. S. deGroot and J. H. van der Waals, *Mol. Phys.*, **6**, 545 (1963).

(4) Nearly quantitative conversions of **1** to **3** were obtained by irradiation of ~0.1% solutions of **1** in a variety of solvents. Compound **3** appears to be identical with that obtained by T. D. Walsh, *J. Am. Chem. Soc.*, **91**, 515 (1969). The authors wish to thank Professor Walsh for providing us with a manuscript of his results prior to publication.

(5) N. Campbell and A. Marks, *J. Chem. Soc.*, 2941 (1951)

(6) J. A. Berson, P. W. Grubb, R. A. Clark, D. R. Hartter, and M. R. Willcott, *J. Am. Chem. Soc.*, **89**, 4076 (1967); J. A. Berson and M. R. Willcott, *ibid.*, **87**, 2751, 2752 (1965); **88**, 2494 (1966); *Record Chem. Progr.* (Kresge-Hooker Sci. Lib.), **27**, 139 (1966).

(7) A. A. Bothner-By, *Advan. Magnetic Resonance*, **1**, 195 (1965).

(8) (a) E. Ciganek, *J. Am. Chem. Soc.*, **87**, 652, 1149 (1965); (b) T. Mukai, H. Kubota, and T. Toda, *Tetrahedron Letters* 3581 (1967), and references therein; (c) W. von E. Doering and M. R. Willcott, *ibid.*, 663 (1962).

(9) Alfred P. Sloan Fellow.

(10) National Institutes of Health Predoctoral Fellow, 1966 to present.

Nicholas J. Turro,⁹ Michael Tobin¹⁰
Department of Chemistry, Columbia University
New York, New York 10027

Lester Friedman, John B. Hamilton
Department of Chemistry, Case Western Reserve University
Cleveland, Ohio 44106
Received November 7, 1968